Algorithm Design and Analysis (XXIII)

MAX-SAT

Guoqiang Li
School of Computer Science

Shanghai Jiao Tong University

MAX-SAT

Given n boolean variables $x_1, \ldots, x_n$, a CNF

$$\varphi(x_1, \ldots, x_n) = \bigwedge_{j=1}^{m} C_j$$

and a nonnegative weight w_j for each C_j.

Find an assignment to the x_is that maximizes the weight of the satisfied clauses.

Simple Randomization Algorithm

Flipping a Coin

Flipping a Coin

A very straightforward randomized approximation algorithm is to set each x_i to `true` independently with probability $1/2$.

Flipping a Coin

A very straightforward randomized approximation algorithm is to set each x_i to `true` independently with probability $1/2$.

Setting each x_i to `true` with probability $1/2$ independently gives a randomized $\frac{1}{2}$-approximation algorithm for weighted MAX-SAT.

Proof

Proof.

Proof

Proof.

Let W be a random variable that is equal to the total weight of the satisfied clauses. Define an indicator random variable Y_j for each clause C_j such that $Y_j = 1$ if and only if C_j is satisfied. Then

$$W = \sum_{j=1}^{m} w_j Y_j$$

We use OPT to denote value of optimum solution, then

$$E[W] = \sum_{j=1}^{m} w_j E[Y_j] = \sum_{j=1}^{m} w_j \cdot \Pr[\text{clause } C_j \text{ satisfied}]$$

Proof (cont'd)

Since each variable is set to `true` independently, we have

$$\Pr[\text{clause } C_j \text{ satisfied}] = \left(1 - \left(\frac{1}{2}\right)^{l_j}\right) \geq \frac{1}{2}$$

where l_j is the number of literals in clause C_j. Hence,

$$E[W] \geq \frac{1}{2}\sum_{j=1}^{m} w_j \geq \frac{1}{2}OPT.$$

A Finer Analysis

A Finer Analysis

Observe that if $l_j \geq k$ for each clause j, then the analysis above shows that the algorithm is a $(1 - (\frac{1}{2})^k)$-approximation algorithm for such instances. For instance, the performance guarantee of MAX E3SAT is $7/8$.

A Finer Analysis

Observe that if $l_j \geq k$ for each clause j, then the analysis above shows that the algorithm is a $(1 - (\frac{1}{2})^k)$-approximation algorithm for such instances. For instance, the performance guarantee of MAX E3SAT is $7/8$.

From the analysis, we can see that the performance of the algorithm is better on instances consisting of long clauses.

A Finer Analysis

Observe that if $l_j \geq k$ for each clause j, then the analysis above shows that the algorithm is a $(1 - (\frac{1}{2})^k)$-approximation algorithm for such instances. For instance, the performance guarantee of MAX E3SAT is $7/8$.

From the analysis, we can see that the performance of the algorithm is better on instances consisting of long clauses.

Theorem

If there is an $(\frac{7}{8} + \epsilon)$-approximation algorithm for MAX E3SAT for any constant $\epsilon > 0$, then **P = NP**.

Derandomization

The previous randomized algorithm can be derandomized. Note that

$$\begin{aligned} E[W] &= E[W \mid x_1 \leftarrow \texttt{true}] \cdot \Pr[x_1 \leftarrow \texttt{true}] \\ &\quad + E[W \mid x_1 \leftarrow \texttt{false}] \cdot \Pr[x_1 \leftarrow \texttt{false}] \\ &= \frac{1}{2}(E[W \mid x_1 \leftarrow \texttt{true}] + E[W \mid x_1 \leftarrow \texttt{false}]) \end{aligned}$$

The previous randomized algorithm can be derandomized. Note that

$$\begin{aligned} E[W] &= E[W \mid x_1 \leftarrow \texttt{true}] \cdot \Pr[x_1 \leftarrow \texttt{true}] \\ &\quad + E[W \mid x_1 \leftarrow \texttt{false}] \cdot \Pr[x_1 \leftarrow \texttt{false}] \\ &= \frac{1}{2}(E[W \mid x_1 \leftarrow \texttt{true}] + E[W \mid x_1 \leftarrow \texttt{false}]) \end{aligned}$$

We set b_1 `true` if $E[W \mid x_1 \leftarrow \texttt{true}] \geq E[W \mid x_1 \leftarrow \texttt{false}]$ and set b_1 `false` otherwise. Let the value of x_1 be b_1.

The previous randomized algorithm can be derandomized. Note that

$$\begin{aligned} E[W] &= E[W \mid x_1 \leftarrow \texttt{true}] \cdot \Pr[x_1 \leftarrow \texttt{true}] \\ &\quad + E[W \mid x_1 \leftarrow \texttt{false}] \cdot \Pr[x_1 \leftarrow \texttt{false}] \\ &= \frac{1}{2}(E[W \mid x_1 \leftarrow \texttt{true}] + E[W \mid x_1 \leftarrow \texttt{false}]) \end{aligned}$$

We set b_1 `true` if $E[W \mid x_1 \leftarrow \texttt{true}] \geq E[W \mid x_1 \leftarrow \texttt{false}]$ and set b_1 `false` otherwise. Let the value of x_1 be b_1.

Continue this process until all b_i are found, i.e., all n variables have been set.

An Example

$$x_3 \vee \overline{x_5} \vee \overline{x_7}$$

An Example

$$x_3 \vee \overline{x_5} \vee \overline{x_7}$$

- $\texttt{Pr}[\text{clause satisfied} \mid x_1 \leftarrow \texttt{true}, x_2 \leftarrow \texttt{false}, x_3 \leftarrow \texttt{true}] = 1$

An Example

$$x_3 \vee \overline{x_5} \vee \overline{x_7}$$

- $\texttt{Pr}[\text{clause satisfied} \mid x_1 \leftarrow \texttt{true}, x_2 \leftarrow \texttt{false}, x_3 \leftarrow \texttt{true}] = 1$
- $\texttt{Pr}[\text{clause satisfied} \mid x_1 \leftarrow \texttt{true}, x_2 \leftarrow \texttt{false}, x_3 \leftarrow \texttt{false}] = 1 - (\frac{1}{2})^2 = \frac{3}{4}$

This is a deterministic $\frac{1}{2}$-approximation algorithm because of the following two facts:

1. $E[W \mid x_1 \leftarrow b_1, \ldots, x_i \leftarrow b_i]$ can be computed in polynomial time for fixed $b_1, \ldots, b_i$.
2. $E[W \mid x_1 \leftarrow b_1, \ldots, x_i \leftarrow b_i, x_{i+1} \leftarrow b_{i+1}] \geq E[W \mid x_1 \leftarrow b_1, \ldots, x_i \leftarrow b_i]$ for all i, and by induction, $E[W \mid x_1 \leftarrow b_1, \ldots, x_i \leftarrow b_i, x_{i+1} \leftarrow b_{i+1}] \geq E[W]$.

Flipping Biased Coins

Flipping Biased Coins

Previously, we set each x_i `true` or `false` with probability $\frac{1}{2}$ independently. $\frac{1}{2}$ is nothing special here.

Flipping Biased Coins

Previously, we set each x_i `true` or `false` with probability $\frac{1}{2}$ independently. $\frac{1}{2}$ is nothing special here.

In the following, we set each x_i `true` with probability $p \geq \frac{1}{2}$.

Flipping Biased Coins

Previously, we set each x_i `true` or `false` with probability $\frac{1}{2}$ independently. $\frac{1}{2}$ is nothing special here.

In the following, we set each x_i `true` with probability $p \geq \frac{1}{2}$.

We first consider the case that no clause is of the form $C_j = \bar{x}_i$.

Flipping Biased Coins

Previously, we set each x_i `true` or `false` with probability $\frac{1}{2}$ independently. $\frac{1}{2}$ is nothing special here.

In the following, we set each x_i `true` with probability $p \geq \frac{1}{2}$.

We first consider the case that no clause is of the form $C_j = \bar{x}_i$.

Lemma

If each x_i is set to `true` with probability $p \geq 1/2$ independently, then the probability that any given clause is satisfied is at least $\min(p, 1 - p^2)$ for instances with no negated unit clauses.

Proof

Proof.

Proof

Proof.

If the clause is a unit clause, then the probability the clause is satisfied is p.

Proof

Proof.

If the clause is a unit clause, then the probability the clause is satisfied is p.

If the clause has length at least two, then the probability that the clause is satisfied is $1 - p^a(1-p)^b$, where a is the number of negated variables and b is the number of unnegated variables. Since $p > \frac{1}{2} > 1 - p$, this probability is at least $1 - p^2$.

Flipping Biased Coins

Armed with previous lemma, we then maximize $\min(p, 1-p^2)$, which is achieved when $p = 1-p^2$, namely $p = \frac{1}{2}(\sqrt{5}-1) \approx 0.618$.

Flipping Biased Coins

Armed with previous lemma, we then maximize $\min(p, 1-p^2)$, which is achieved when $p = 1-p^2$, namely $p = \frac{1}{2}(\sqrt{5}-1) \approx 0.618$.

We need more effort to deal with negated unit clauses, i.e., $C_j = \bar{x}_i$ for some j.

Armed with previous lemma, we then maximize $\min(p, 1-p^2)$, which is achieved when $p = 1-p^2$, namely $p = \frac{1}{2}(\sqrt{5}-1) \approx 0.618$.

We need more effort to deal with negated unit clauses, i.e., $C_j = \bar{x}_i$ for some j.

We distinguish between two cases:

1. Assume $C_j = \bar{x}_i$ and there is no clause such that $C = x_i$. In this case, we can introduce a new variable y and replace the appearance of $\bar{x}_i$ in φ by y and the appearance of x_i by $\bar{y}$.

2. $C_j = \bar{x}_i$ and some clause $C_k = x_i$. W.L.O.G we assume $w(C_j) \leq w(C_k)$. Note that for any assignment, C_j and C_k cannot be satisfied simultaneously. Let v_i be the weight of the unit clause $\bar{x}_i$ if it exists in the instance, and let v_i be zero otherwise, we have

$$\mathrm{OPT} \leq \sum_{j=1}^{m} w_j - \sum_{i=1}^{n} v_i$$

2. $C_j = \bar{x}_i$ and some clause $C_k = x_i$. W.L.O.G we assume $w(C_j) \leq w(C_k)$. Note that for any assignment, C_j and C_k cannot be satisfied simultaneously. Let v_i be the weight of the unit clause $\bar{x}_i$ if it exists in the instance, and let v_i be zero otherwise, we have

$$\mathrm{OPT} \leq \sum_{j=1}^{m} w_j - \sum_{i=1}^{n} v_i$$

We set each x_i `true` with probability $p = \frac{1}{2}(\sqrt{5} - 1)$, then

$$\begin{aligned} E[W] &= \sum_{j=1}^{m} w_j E[Y_j] \\ &\geq p \cdot \left(\sum_{j=1}^{m} w_j - \sum_{i=1}^{n} v_i \right) \\ &\geq p \cdot \mathrm{OPT} \end{aligned}$$

Rounding by Linear Programming

The Use of Linear Program

Integer Program Characterization:

$$
\begin{array}{lll}
\max & \sum_{j=1}^{m} w_j z_j & \\
& \sum_{i \in P_j} y_i + \sum_{i \in N_j} (1 - y_i) \geq z_j, & \forall C_j = \bigvee_{i \in P_j} x_i \vee \bigvee_{i \in N_j} \bar{x}_i, \\
& y_i \in \{0, 1\}, & i = 1, \ldots, n, \\
& z_j \in \{0, 1\}, & j = 1, \ldots, m.
\end{array}
$$

where y_i indicate the assignment of variable x_i and z_j indicates whether clause C_j is satisfied.

The Use of Linear Program

Linear Program Relaxation:

$$
\begin{array}{lll}
\max & \sum_{j=1}^{m} w_j z_j & \\
& \sum_{i \in P_j} y_i + \sum_{i \in N_j} (1 - y_i) \geq z_j, & \forall C_j = \bigvee_{i \in P_j} x_i \vee \bigvee_{i \in N_j} \bar{x}_i, \\
& 0 \leq y_i \leq 1, & i = 1, \ldots, n, \\
& 0 \leq z_j \leq 1, & j = 1, \ldots, m.
\end{array}
$$

where y_i indicate the assignment of variable x_i and z_j indicates whether clause C_j is satisfied.

Flipping Different Coins

Let (y^*, z^*) be an optimal solution of the linear program.

Flipping Different Coins

Let (y^*, z^*) be an optimal solution of the linear program.

We set x_i to `true` with probability y_i^*.

Flipping Different Coins

Let (y^*, z^*) be an optimal solution of the linear program.

We set x_i to `true` with probability y_i^*.

This can be viewed as flipping different coins for every variable.

Flipping Different Coins

Let (y^*, z^*) be an optimal solution of the linear program.

We set x_i to `true` with probability y_i^*.

This can be viewed as flipping different coins for every variable.

Theorem

Randomized rounding gives a randomized $(1 - \frac{1}{e})$-approximation algorithm for MAX-SAT.

Analysis

$$
\begin{aligned}
& \Pr[\text{clause } C_j \text{ not satisfied}] \\
= \quad & \prod_{i \in P_j} (1 - y_i^*) \prod_{i \in N_j} y_i^* \\
\leq \quad & \left[\tfrac{1}{l_j} \left(\textstyle\sum_{i \in P_j} (1 - y_i^*) + \sum_{i \in N_j} y_i^* \right) \right]^{l_j} \\
= \quad & \left[1 - \frac{1}{l_j} \left(\sum_{i \in P_j} y_i^* + \sum_{i \in N_j} (1 - y_i^*) \right) \right]^{l_j} \leq \left(1 - \frac{{z_j^*}^{l_j}}{l_j} \right)
\end{aligned}
$$

Analysis

$$
\begin{aligned}
& \Pr[\text{clause } C_j \text{ not satisfied}] \\
= \quad & \prod_{i \in P_j} (1 - y_i^*) \prod_{i \in N_j} y_i^* \\
\leq \quad & \left[\tfrac{1}{l_j} \left(\textstyle\sum_{i \in P_j} (1 - y_i^*) + \sum_{i \in N_j} y_i^* \right) \right]^{l_j} \\
= \quad & \left[1 - \frac{1}{l_j} \left(\sum_{i \in P_j} y_i^* + \sum_{i \in N_j} (1 - y_i^*) \right) \right]^{l_j} \leq \left(1 - \frac{{z_j^*}^{l_j}}{l_j} \right)
\end{aligned}
$$

Arithmetic-Geometric Mean Inequality

Analysis

$$
\begin{aligned}
& \Pr[\text{clause } C_j \text{ satisfied}] \\
\geq\ & 1 - \left(1 - \frac{z_j^*}{l_j}\right)^{l_j} \\
\geq\ & \left[1 - \left(1 - \tfrac{1}{l_j}\right)^{l_j}\right] z_j^*
\end{aligned}
$$

Analysis

$$
\begin{aligned}
& \Pr[\text{clause } C_j \text{ satisfied}] \\
\geq\ & 1 - \left(1 - \frac{z_j^*}{l_j}\right)^{l_j} \\
\geq\ & \left[1 - \left(1 - \tfrac{1}{l_j}\right)^{l_j}\right] z_j^* \qquad \text{Jensen's Inequality}
\end{aligned}
$$

Analysis

$$
\begin{aligned}
& \Pr[\text{clause } C_j \text{ satisfied}] \\
\geq\ & 1 - \left(1 - \frac{z_j^*}{l_j}\right)^{l_j} \\
\geq\ & \left[1 - \left(1 - \tfrac{1}{l_j}\right)^{l_j}\right] z_j^* \qquad \text{Jensen's Inequality}
\end{aligned}
$$

Therefore, we have

$$
\begin{aligned}
E[W] &= \sum_{j=1}^{m} w_j \Pr[\text{clause } C_j \text{ satisfied}] \\
&\geq \sum_{j=1}^{m} w_j z_j^* \left[1 - \left(1 - \frac{1}{l_j}\right)^{l_j}\right] \\
&\geq \left(1 - \frac{1}{e}\right) \cdot \mathrm{OPT}
\end{aligned}
$$

The Combined Algorithm

Choosing the Better of Two

The randomized rounding algorithm performs better when l_j-s are small. ($\left(1-\frac{1}{k}\right)^k$ is nondecreasing)

Choosing the Better of Two

The randomized rounding algorithm performs better when l_j-s are small. ($\left(1-\frac{1}{k}\right)^k$ is nondecreasing)

The unbiased randomized algorithm performs better when l_j-s are large.

Choosing the Better of Two

The randomized rounding algorithm performs better when l_j-s are small. ($\left(1-\frac{1}{k}\right)^k$ is nondecreasing)

The unbiased randomized algorithm performs better when l_j-s are large.

We will combine them together.

Choosing the Better of Two

The randomized rounding algorithm performs better when l_j-s are small. ($\left(1-\frac{1}{k}\right)^k$ is nondecreasing)

The unbiased randomized algorithm performs better when l_j-s are large.

We will combine them together.

Theorem

Choosing the better of the two solutions given by the two algorithms yields a randomized $\frac{3}{4}$-approximation algorithm for MAX-SAT.

Analysis

Let W_1 and W_2 be the r.v. of value of solution of randomized rounding algorithm and unbiased randomized algorithm respectively. Then

$$\begin{aligned}
E[\max(W_1, W_2)] &\geq E[\frac{1}{2}W_1 + \frac{1}{2}W_2] \\
&\geq \frac{1}{2}\sum_{j=1}^{m} w_j z_j^* \left[1 - \left(1 - \frac{1}{l_j}\right)^{l_j}\right] + \frac{1}{2}\sum_{j=1}^{m} w_j \left(1 - 2^{-l_j}\right) \\
&\geq \sum_{j=1}^{m} w_j z_j^* \left[\frac{1}{2}\left(1 - \left(1 - \frac{1}{l_j}\right)^{l_j}\right) + \frac{1}{2}\left(1 - 2^{-l_j}\right)\right] \\
&\geq \frac{3}{4} \cdot \mathrm{OPT}
\end{aligned}$$

Referred Materials

Referred Materials

Content of this lecture comes from Chapter 16 in [Vaz04].